

REGLAN DRUG-BLACK BOX WARNING!

“On February 26, 2009 the U.S. Food & Drug Administration (FDA) added a “Black Box Warning” to metoclopramide - containing drugs including Reglan tablets, Reglan oral disintegrating tablet, metoclopramide oral solution, and Reglan injections.”



REGLAN is a drug to treat gastrointestinal disorders. Chronic use of *Reglan* has been linked to *Tardive Dyskinesia*, which may include involuntary and repetitive movements of the body, even after the drugs are no longer taken. The development of this condition is directly related to the length of time a patient is taking *Reglan* and the number of doses taken. Those at greatest risk include the elderly, especially older women, children and people who have been on the drug for a long time.

TARDIVE DYSKINESIA - is a serious and often irreversible movement disorder with the following characteristics: Repetitive movement of extremities, facial ticks, constant lip smacking, unintentional grimacing of the face, tongue protrusion, rapid eye movement or blinking, puckering of lips, and impaired finger movement.

People suffering from *Tardive Dyskinesia* often withdraw from social contact, and can be effected emotionally and psychologically - leading to severe depression.

If you or a loved one has taken *Reglan* and has been diagnosed with *Tardive Dyskinesia*, contact **BLACKMON & BLACKMON, PLLC** to learn how we can help you recover financially from any temporary or permanent side effects related to this dangerous drug. Please contact us by calling **1-888-858-1567 or (601) 859-1567.**